

## INSIDE THIS ISSUE:

Compact Fluorescents	1
Fragrance: Carcinogenic	9
Teaching Medical Prof.	10
Science Lacks Integrity	12
Children in Danger	15
Money Spin- ning Fungi	17
Violence Not Behavioral	18
Fragrances Toxic	19
Hospital Accessibility	20
Fibromyalgia and ATP	25
Disability: A Free Ride?	26
Patient Resources	28
Community News	33
Research Studies	34

# MCSA NEWS

VOLUME 3

ISSUE 8

AUGUST 2008

## Compact Fluorescent Lamps (CFLs)



## What You Need to Know About Low Energy Lighting

# Compact Fluorescent Lamps

## *What You Need to Know About Low Energy Lighting*

*-Dr. Andrew Goldsworthy*

**“The  
principle of  
operation  
is the same  
as a  
fluorescent  
strip-light.”**

### **Introduction**

Compact fluorescent lamps (CFLs) are smaller versions of the familiar fluorescent strip-lights found in schools, public buildings and many people's kitchens. Like the strip-lights, they are about five times more efficient than tungsten incandescent lamps at turning electrical power into light. They also last many times longer and the saving in energy over their lifetime more than offsets their extra cost. Governments all over the world are either encouraging or coercing us to replace our tungsten lamps with CFLs to save energy and reduce our carbon footprint.

The principle of operation is the same as a fluorescent strip-light. An electric current is driven through a tube containing argon and a small amount of mercury vapor. This generates invisible ultra-violet light that excites a fluorescent coating (the phosphor) on the inside of the tube, which then emits longer-wavelength visible light.

### **Environmental impact**

Unlike incandescent lamps, CFLs contain toxic chemicals. Each one contains about 4mg of mercury, which is a cumulative poison. However, because coal also contains mercury, which is released into the atmos-

phere when burned, this too is a source of mercury pollution. If we assume that all our electricity came from coal, then the amount of mercury pollution saved by switching to CFLs is about double that in the lamps themselves

<http://tinyurl.com/68t3u6>

so their use could reduce the net mercury burden on the environment.

Nevertheless, there can be problems with local pollution if they are not properly disposed of. In Europe, there are regulations requiring retailers of CFLs to provide free facilities for their recycling, but these are poorly implemented in the UK. Most of them still end up in land-fill, where they may be broken and release their mercury and other toxins. This can give high local concentrations, with a risk of contamination to water supplies.

### **Breakage**

We also have to think of what to do if we actually break one indoors. Because mercury vapor is toxic, the best solution is to open the windows and vacate the room for about 15 minutes until the mercury vapor clears. Then wear rubber gloves to clear up the fragments (which also contain toxic phosphors) with a dustpan and brush (not a vacuum cleaner). Any remaining shards of glass should be cleaned up with a moist paper towel and everything double bagged for disposal.



**“Most  
of  
them  
cannot  
be  
dimmed  
with  
dimmer  
switches”**



### **Light output**

CFLs are physically larger than the equivalent tungsten lamps and you may have to use a smaller and dimmer one if it is to fit into an existing fitting. They are also not best suited for outdoor use since they perform poorly in the cold. Even indoors, many of them can take several minutes to reach full brightness and are unsuitable for short periods of use such as in a toilet. Not only may they not reach full brightness during your visit, but their life span will probably be reduced to no more than that of an incandescent lamp under these conditions. A further problem with their brightness is that most of them cannot be dimmed with dimmer switches since they tend to be either fully on or fully off.

### **Color of the light**

The color of a fluorescent lamp is usually described by its color temperature, which is the temperature to which a metal would have to be heated to give that color. For example, a warm white lamp has a color temperature of around 2700 degrees Kelvin (Celsius + 273) whereas natural noon daylight is somewhere between 5000 and 6000 degrees Kelvin. Different colors are obtained by choosing different phosphors. Often, there is a mixture of phosphors to give something that looks like daylight. However, this is an illusion. Real daylight consists of a broad spectrum of all wavelengths, but fluorescent light is a mixture of peaks at different wavelengths with dark areas in between.

### **Color-rendering**

Even a "daylight" fluorescent lamp doesn't give the equivalent of true daylight

because of the gaps in its spectrum. These gaps reduce the "richness" of the colors seen under its light and it makes accurate color matching difficult. It is possible to fill some of the gaps by adding extra phosphors, but these also reduce the efficiency of the lamp so that the number normally added is a compromise. Just how good a particular lamp is for matching colors is measured as its color-rendering index. A continuous spectrum from daylight or a tungsten lamp is taken as 100, whereas a fluorescent lamp may have a color-rendering index of between about 50 (very poor) and over 90 (good). Triphosphor lamps give good but not perfect, color rendering with a near daylight color temperature. However, many people who are used to incandescent lighting find them too "cold" for a living room and prefer the warmer colors such as warm white.

### **Electronics**

Fluorescent lamps will only run on alternating current. They also need a pulse of high voltage and heated filaments at either end to start the electrical discharge that lights them. After that, the current must be limited externally, otherwise too much would flow and they would burn out. In a traditional fluorescent strip light, this is accomplished by the starter switch and the choke (a coil of wire wound around an iron core). Once started, the current flows through the tube as a smooth sine wave at mains frequency, which is 50Hz (cycles per second) in Europe and 60Hz in America. This makes the light flash on and off with each half cycle (i.e. 100 or 120 times a second) and some people, such as epileptics and migraine sufferers find this disturbing.

**“Many  
people  
have  
reported  
ill effects  
when  
using  
CFLs.”**



However, almost all CFLs use electronic control gear. This usually incorporates a switched-mode power supply in the base of the lamp itself. It rectifies the AC from the mains to convert it to DC and then chops it electronically into a series of sharp rectangular alternating pulses, which then light the lamp. However, the new frequency, which is usually about 40 kHz (40,000 cycles per second), is so high and the gaps between pulses are so short that the relatively slow response of the phosphors can fill them easily. Consequently, these lamps do not flash.

#### **Biological effects**

Despite the absence of flashing, many people have reported ill effects when using CFLs. Typical symptoms include dizziness, nausea, tinnitus (ringing or buzzing in the ears), headaches and various skin disorders. In particular, many sufferers from migraine and epilepsy have found that they still aggravate their conditions.

<http://tinyurl.com/6449ed>

<http://tinyurl.com/6c6hnz>

The effects may be due to pulsed electromagnetic radiation.

The symptoms of exposure to CFL radiation are remarkably similar to those reported by electrosensitive individuals when exposed to pulsed electromagnetic fields. Since the lamps do not flash, it seems probable that they are a direct effect of the pulsed radiation on the brain and nervous system. The magnetic component of the radiation is the more dangerous because it can penetrate deep into the human body where it generates electrical voltages proportional to its rate of change. The rapid rise and fall times of these magnetic pulses can therefore give relatively massive and po-

tentially damaging voltage spikes both in living cells and across their membranes.

#### **Contamination of the mains**

Poor quality CFLs often allow these pulses to leak back into the mains wiring to contribute to "dirty electricity" and increase the range of their effects to neighboring rooms or houses. You should be able to detect these by holding a portable radio tuned between stations on an AM band near the wiring. This is because pulses, by their very nature, also contain harmonics (multiples of the original frequency) that can extend well into the radio frequency spectrum. If you hear a buzzing sound from the set, it means that pulses are leaking into the mains and you should replace the offending lamp by another of better quality.

Contamination of the mains to give "dirty electricity" can come from many sources, not just CFLs. Measurements made by David Stetzer in the library of an American school showed it to consist of hundreds of sharp spikes that could be up to hundreds of millivolts high, superimposed on each cycle of the 120 volt mains supply. Although the largest of them was only a tiny fraction of the overall mains voltage, their rapid rise and fall times give them biological activity. The sharp magnetic spikes they generate penetrate living tissue easily, where their sudden changes in field-strength induce large voltage spikes. Several studies by Dr Magda Havas of Trent University in Canada and various co-workers have shown that simply removing these spikes in the mains with "Graham/Stetzer" filters gave improvements in the health, learning ability and behavior of schoolchildren, reductions in the insulin needed to treat diabetics and an alleviation of the symptoms of electrosensitivity.

**“Only about three percent of the population are thought to suffer from electromagnetic hypersensitivity.”**

### **Electrosensitivity**

People who are affected badly by weak electromagnetic fields in this way are described as being electrosensitive or as suffering from electromagnetic hypersensitivity (EHS). Only about three percent of the population are thought to suffer from EHS at present, although this proportion is expected to rise as more people become sensitized and people who are already sensitive but do not realize it discover that their symptoms are related to electromagnetic exposure.

The symptoms of electrosensitivity are many and varied and not everyone suffers in the same way or to the same degree. Some of the effects are on the brain and nervous system and often become apparent during or shortly after exposure. They include dizziness, tinnitus, pins and needles, sensations of burning, numbness, fatigue and headaches. Longer-term effects include skin disorders, gut problems and an increased tendency to allergies and multiple chemical sensitivities.

See <http://tinyurl.com/6qs8ew>.

### **Mechanisms of electrosensitivity**

Electrosensitive individuals are physiologically different to the rest of the community. Eltiti and her co-workers at Essex University showed this very clearly in a project for the mo-

bile phone industry and the UK Government. They wanted to see if electrosensitive individuals could detect the radiation from mobile phone masts. They excluded epileptics and people wearing pacemakers for cardiac arrhythmia who might be particularly sensitive and most of their results were less conclusive than they should have been. However, they did show very clearly that their group of EHS sufferers had skins with a significantly higher electrical conductance than the non-sensitive controls ( $p < 0.001$ ). This means that their skin cells were more permeable to ions (charged atoms and molecules) that normally carry electricity in living tissues. There is now considerable evidence that most of the symptoms of electrosensitivity result from ions leaking through membranes in response to electromagnetic fields. Consequently, if electrosensitive individuals already have abnormally leaky membranes, they will be more likely to be affected by these fields.

### **Sensory disturbances**

Membrane leakage can account for the neurological symptoms of EHS sufferers. We know that weak electromagnetic radiation can temporarily remove structurally important calcium ions from cell membranes to make them leak.

<http://tinyurl.com/2nfujj>

Unfortunately, all of our senses depend on ions flowing through the membranes of sensory cells at a rate that depends on the strength of the stimulus. This works well for most of us most of the time, but if the sensory cells of electrosensitive individuals are already leaky, any further electromagnetically-induced leakage will be more likely to trigger them to generate nerve impulses and give false sensations.



**“Electromagnetic radiation can in some people cause tinnitus, which can range from a mild ringing in the ears to buzzing.”**

### **The effects on the ear are like motion sickness**

The main sensory cells of the ear are the hair cells. Hairs at the apices of these bend when they sense movement in the surrounding medium. This makes ions leak through their membranes to reduce the voltage across them. They respond by releasing neurotransmitters that stimulate neighboring nerve cells to send signals to the brain. Those at the ends of the semicircular canals have their hairs embedded in a light jelly, which deforms in response to movements of the fluid within. Because the fluid inside the canals tends to stay stationary when the head twists suddenly, it appears to flow past the jelly so that it measures rapid changes in the orientation of the head. The jelly in other parts of the ear is weighted with mineral granules (otoliths) and deforms in response to gravity and linear acceleration. The hair cells in these regions act like plumb-lines and give us most of our sense of balance.

We are all familiar with what happens if we feed them false information. If we spin our bodies rapidly and suddenly stop, the fluid in the semicircular canals continues to swirl for a while, the signals from the hair cells conflict with what we see around us and we feel dizzy. The stress and nausea of people who get motion sickness is due to a similar conflict between the signals from the ear and those from the other senses such as touch, sight and pressure on specific regions of the skin. It is therefore not surprising that false signals generated by electromagnetically-induced leakage in the hair cells cause dizziness and nausea in some

electrosensitive individuals.

### **It can also cause tinnitus**

The hair cells in the cochlea (the hearing part of the inner ear) respond to sound. They are arranged in a graded sequence with different length hairs along the length of the cochlea. Like the strings of a harp, they resonate at different frequencies. When an incoming sound matches their resonant frequencies, the hairs vibrate more strongly. This makes the cells concerned leak more ions, and trigger neighboring nerve cells to send impulses to the brain. Which cells are stimulated tell it the pitch of the note. The frequency of the impulses tells it the loudness. False stimulation of these cells by electromagnetic radiation can in some people cause tinnitus, which can range from a mild ringing in the ears to buzzing and complex sounds that may be loud enough to drown out normal conversation.

### **Effects on the other senses**

There are countless cells all over our bodies that sense various forms of touch (mechanoreceptors) temperature (thermoreceptors) and pain (nociceptors). Each group contains many specialized variants but they nearly all function by letting ions flow through their membranes at a rate that depends on the strength of the stimulus. This reduces the voltage across the cell membrane, which triggers the transmission of nerve impulses to the brain, either by the cell itself or by releasing neurotransmitters to stimulate neighboring nerve cells. Electromagnetically-induced membrane leakage in sensory cells in the skin explains the pins and needles, sensations of burning and pain experienced by EHS sufferers.



**“False  
nerve  
impulses  
can be  
generated  
by  
electromagnetic  
fields in the  
neurons  
of the brain.”**

### **The eye is different**

The light-sensing rods and cones in the retina of the eye are an exception in that when they respond to light they increase rather than decrease the voltage across their membranes. Consequently, any uncontrolled electromagnetically induced leakage here might be expected to reduce their sensitivity. It may be no coincidence that electrosensitive people whose vision is affected usually report a blurring or partial loss of vision rather than seeing things that aren't there.

### **Effects on the brain**

It isn't just the sensory cells that are affected by electromagnetic radiation. False nerve impulses can be generated by electromagnetic fields in the neurons of the brain. These can cause hyperactivity, make it more difficult to sleep, trigger random thoughts, and result in a loss of concentration and confused thinking.

<http://tinyurl.com/55286a>

four times more likely to have an accident by using a mobile phone when driving, since using a hands-free type is no better but talking to a passenger has little or no effect.

### **Non-neurological effects**

Spurious action potentials caused by membrane leakage in the heart muscle can give rise to cardiac arrhythmia and an increased risk of heart attacks. Increases in the permeability of skin cells can give rise to dermatological problems as well as a greater tendency to develop allergies and multiple chemical sensitivities. Electromagnetically-induced increases in the permeability of the gut to toxins, carcinogens and its partially digested contents might be expected to cause a whole array of disorders and have been implicated as a risk factor in the development of autoimmune diseases such as multiple sclerosis and type-I diabetes.

<http://tinyurl.com/55286a>

All of these illnesses have been linked scientifically to electromagnetic exposure, so people with a tendency to any of them should take the utmost caution in the use of CFLs and avoid using them totally if possible.

### **Are there alternatives?**

If you are affected by CFLs, an obvious solution is to stock up on incandescent bulbs before they are phased out. If this is not an option, try using high voltage halogen incandescent lamps as a replacement since there are no immediate plans to phase these out. However, do not use the low voltage types, since many of them use switched mode power supplies to reduce the voltage. These could well give the same symptoms as CFLs.



It may therefore not be advisable to use CFLs in a study or any other place where a great deal of concentration is required, especially if you are electrosensitive. This effect is probably the real reason why we are

**“CFLs are  
not  
the best  
option  
for low  
energy  
lighting.”**

### **What next?**

It is becoming increasingly obvious that CFLs are not the best option for low energy lighting, and special dispensation needs to be made to supply alternatives to people whose health is unduly affected by them. Even so, we should regard CFLs as being just a stopgap until LED (light emitting diode) lighting is perfected. LEDs last indefinitely, they run on DC or rectified AC without generating damaging electromagnetic pulses, and the best of them are already more efficient than CFLs. At the moment, the main problem with them is with their color; the most efficient "white" ones have a harsh blue tint. Although they are commonly used in flashlights, they have very poor color

rendering abilities and aren't really suitable for domestic lighting. Their spectrum can be improved by adding phosphors to absorb some of the blue light and re-emit it as other colors, but this causes a dramatic loss of efficiency. An alternative is to use an array of differently colored LEDs so that between them they give a spectrum that corresponds more closely to true white light. Hopefully, research on these devices will be given a high priority so that cheap high-quality LED lighting for domestic and industrial installations becomes available and CFLs, with all their attendant problems, become things of the past.

Copyrighted © 2008  
Dr. Andrew Goldsworthy  
Reprinted with Permission



Andrew Goldsworthy was born in 1939. After a conventional Grammar School education he obtained a First Class Honours Degree in Botany followed by a PhD for research into plant physiology and biochemistry at the University of Wales. He went on to lecture at Imperial College London, where he spent the rest of his career. He has had many teaching and research interests, ranging from the biochemistry of photorespiration to the biology of space flight. He retired in 2004 but remains as an honorary lecturer, and still gives occasional lectures on specialized subjects. He was also a scientific advisor to the European Space Agency and is currently a scientific advisor to several European charities concerned with the environment and electromagnetic fields, including the h.e.s.e - project, the Radiation Research Trust and Electrosensitivity-UK. He has always had a strong interest in how living organisms use internally-generated electric currents to control their growth and metabolism and in their disruption by externally-applied currents and fields. In his retirement, he pieced together nuggets of information from a wide range of scientific journals and created a simple layperson's explanation of how weak electromagnetic fields affect us all. This can be found on the h.e.s.e. - project website at:

<http://tinyurl.com/28lo82>,

which corresponds to:

<http://www.hese-project.org/hese-uk/en/niemr/resonance1.php>.